

The mean perimeter of some random plane convex sets generated by a Brownian motion

Philippe Biane*

Gérard Letac†

Abstract

If C_1 is the convex hull of the curve of the standard Brownian motion in the complex plane watched from 0 to 1, we consider the convex hulls of C_1 and several rotations of it and we compute the mean of the length of their perimeter by elementary calculations.

I Introduction

If C is a compact plane convex set, denote by $L(C)$ the length of its perimeter. Let $B(t) = (X(t), Y(t))$ be a standard Brownian motion in the plane and denote by $C_1(t)$ the convex hull of the curve

$$\mathbf{B} = \{B(s) ; 0 \leq s \leq t\}.$$

This object has been considered by Paul Lévy (1948), pages 239-240. The first author a long time ago has called attention on the fact that the mean of $L(C_1(t))$ is $\sqrt{8\pi t}$ in Letac (1978). Here we consider the symmetric convex hull $C_2(t)$, namely the convex hull of $C_1(t) \cup -C_1(t)$. The convex set $C_2(t)$ should not be confused with the Minkowski sum $C_1(t) + (-C_1(t))$, which is bigger. By scaling we can guess the existence of a constant ℓ_2 such that $E[L(C_2(t))] = \ell_2\sqrt{8\pi t}$. Similarly we consider random convex sets obtained in the following way. We let $\Omega \subset [0, 2\pi[$ be a set of angles and C_Ω the convex hull of $\cup_{\omega \in \Omega} R_\omega \mathbf{B}$, where R_ω is the rotation of angle ω . Again by scaling there exists a constant ℓ_Ω such that $E[L(C_\Omega(t))] = \ell_\Omega\sqrt{8\pi t}$. We shall compute these constants when Ω is equal to one of the following sets

$$\{0\}, \quad \{0, \pi\}, \quad \{0, \pi/2\}, \quad \{0, 2\pi/3\}, \quad \{0, \pi/2, \pi\}$$

or

$$\{0, 2\pi/3, 4\pi/3\} \quad \{0, \pi/2, \pi, 3\pi/2\}, \quad [0, 2\pi[$$

giving the result either in closed form or as an expression which can be evaluated numerically.

*CNRS, Laboratoire d'Informatique, Institut Gaspard Monge, Université Paris-Est, 5 Boulevard Descartes, Champs-sur-Marne F-77454, Marne-la-Vallée Cedex 2 Philippe.Biane@univ-mlv.fr

†Laboratoire de Statistique et Probabilités, Université Paul Sabatier, 31062, Toulouse, France, letac@cict.fr

II Computations

II.1 Preliminary remarks

Given a compact convex set C in the Euclidean plane $\mathbb{R}^2$ consider its support function h_C as the function on $\mathbb{R}$ of period 2π defined by

$$h_C(\theta) = \max_{(x,y) \in C} (x \cos \theta + y \sin \theta).$$

It is well known (see Letac (1983), formula (7.2) for a proof without regularity hypothesis) that the length of the perimeter of C is $\int_0^{2\pi} h_C(\theta) d\theta$. This formula is generally attributed to Cauchy. Let $\Omega \subset [0, 2\pi[$ and define $h_{\mathbf{B}}(\theta) = \max_{0 \leq s \leq t} (X(s) \cos \theta + Y(s) \sin \theta)$ and

$$h_{\Omega}(\theta) = \sup_{\omega \in \Omega} h_{\mathbf{B}}(\theta + \omega),$$

then rotation invariance of the law of Brownian motion implies

$$E[L(C_{\Omega}(t))] = 2\pi E[h_{\Omega}(0)].$$

Let P_{Ω} be the convex set defined by the inequalities

$$x \cos \omega + y \sin \omega \leq 1; \omega \in \Omega.$$

In other terms, P_{Ω} is the intersection of half planes whose boundary is tangent to the unit circle. Denote by T_{Ω} the first time where Brownian motion exits this set. Then a scaling argument shows that $h_{\Omega}(0)$ is distributed as $T_{\Omega}^{-1/2}$. It turns out that in several cases the Laplace transform $E[\exp(-\frac{\lambda^2}{2} T_{\Omega})]$ is given by a simple formula. Under such a circumstance

$$E[h_{\Omega}(0)] = E[T_{\Omega}^{-1/2}] = \sqrt{\frac{2}{\pi}} \int_0^{\infty} E[\exp(-\frac{\lambda^2}{2} T_{\Omega})] d\lambda$$

and

$$E[L(C_{\Omega})(t)] = \sqrt{8\pi t} \int_0^{\infty} E[\exp(-\frac{\lambda^2}{2} T_{\Omega})] d\lambda. \quad (2.1)$$

II.2 Some easy cases

II.2.1 $\Omega = \{0\}$

The time T_{Ω} is the first hitting time of 1 for a linear Brownian motion, and

$$E[\exp(-\frac{\lambda^2}{2} T_{\Omega})] = e^{-\lambda}. \quad (2.2)$$

The Laplace tranform can be inverted to give

$$P(T_{\Omega} \in dt) = \frac{e^{-1/2t}}{\sqrt{2\pi t^3}}. dt \quad (2.3)$$

Using (2.1) we get $E[L(C_1(t))] = \sqrt{8\pi t}$. As expected: $l_{\Omega} = 1$.

II.2.2 $\Omega = \{0, \pi\}$

Now T_Ω is the exit time of linear Brownian motion from the interval $[-1, +1]$ and

$$E[\exp(-\frac{\lambda^2}{2}T_\Omega)] = 1/\cosh \lambda \quad (2.4)$$

with density

$$P(T_\Omega \in dt) = \pi \sum_{n=0}^{\infty} (-1)^n (n + \frac{1}{2}) e^{-(n+\frac{1}{2})^2 \pi^2 t/2} dt \quad (2.5)$$

and distribution function

$$P(T_\Omega \geq t) = \frac{2}{\pi} \sum_{n=0}^{\infty} \frac{(-1)^n}{n + \frac{1}{2}} e^{-(n+\frac{1}{2})^2 \pi^2 t/2} \quad (2.6)$$

see e.g. Biane, Pitman, Yor (2001). Using again (2.1) gives

$$E[L(C_2(t))] = \frac{\pi}{2} \sqrt{8\pi t}.$$

Thus $\ell_\Omega = \frac{\pi}{2}$.

II.2.3 $\Omega = [0, 2\pi[$

The set P_Ω is the unit circle, and the time T_Ω is the first hitting time of 1 by a Bessel process of dimension 2, therefore

$$\mathbb{E}(e^{-\frac{\lambda^2}{2}T_1}) = \frac{I_0(0)}{I_0(\lambda)} \quad (2.7)$$

where

$$I_0(2z) = \sum_{k=0}^{\infty} \frac{z^{2k}}{k!^2}$$

is the Bessel function. Thus

$$\ell_\Omega = \int_0^\infty \frac{d\lambda}{I_0(\lambda)}.$$

A calculation by Mathematica (thanks to Daoud Bshouty for that) gives $\ell_\Omega = 2,08323..$

II.3 A cone of angle $\pi/3$

We consider the case $\Omega = \{0, 2\pi/3\}$. Therefore T_Ω is the exit time from a cone of angle $\pi/3$ tangent to the unit circle. In this case one can use results from Doumerc and O'Connell (2005). There, the exit time from the cone $x_1 > x_2 > x_3$ in $\mathbf{R}^3$, starting from the point (x_1, x_2, x_3) is expressed as

$$P(T \geq t) = p_{12} - p_{13} + p_{23}$$

with $p_{ij} = \sqrt{\frac{2}{\pi}} \int_0^{(x_i - x_j)/\sqrt{2t}} e^{-y^2/2} dy$. Taking the orthogonal projection of Brownian motion on the hyperplane $x_1 + x_2 + x_3 = 0$, with the starting point $(\sqrt{2}, 0, -\sqrt{2})$, one gets

$$P(T_\Omega \geq t) = \sqrt{\frac{2}{\pi}} \left(2 \int_0^{1/\sqrt{t}} e^{-y^2/2} dy - \int_0^{2/\sqrt{t}} e^{-y^2/2} dy \right)$$

and a straightforward computation gives

$$E[T_\Omega^{-1/2}] = \frac{3}{2} \sqrt{\frac{2}{\pi}}$$

and

$$E[L(C_\Omega(t))] = \frac{3}{2} \sqrt{8\pi t}.$$

Thus $\ell_\Omega = 3/2$.

II.4 An equilateral triangle

An interesting case is when $\Omega = \{0, 2\pi/3, 4\pi/3\}$ and P_Ω is an equilateral triangle centered at 0. We will compute the distribution of T_Ω and get a curious identity in law.

In order to do the computation it will be convenient to take the equilateral triangle Δ with vertices $0, 1, -j^2$ where $j = (-1 + i\sqrt{3})/2$. Its center is $x_0 = (1 - j^2)/3$. The case of the triangle P_Ω can be obtained by an affinity. The orthogonal reflections with respect to the three lines bordering the triangle Δ generate a group W of affine isometries of the Euclidian plane, whose fundamental domain is Δ . Using the reflection principle, we can compute the probability transition of Brownian motion in the triangle, killed at the boundary. One obtains

$$p_t^0(x, y) = \sum_{w \in W} \det(w) p_t(x, w(y))$$

where $p_t(x, y)$ is the ordinary heat kernel on the Euclidean plane and x and y are points of this plane. If we denote R the lattice generated by 1 and j , then the dual lattice $\hat{R}$ (such that $\langle w, \hat{w} \rangle \in \mathbf{Z}$) is generated by the dual basis $a = 1 + \frac{i}{\sqrt{3}}$ and $b = \frac{2i}{\sqrt{3}}$. The plane is tiled by translates of Δ by R , which correspond to direct isometries $w \in W$ and translates of $\bar{\Delta}$. Thus

$$p_t^0(x, y) = \sum_{r \in R} p_t(x - y, r) - p_t(x - \bar{y}, r).$$

The function

$$h(t, z) = \sum_{r \in R} p_t(z, r)$$

is periodic in z with period lattice R and $p_t^0(x, y) = h(t, x - y) - h(t, x - \bar{y})$. One can apply Poisson summation formula, i.e. expand it in terms of the characters $e^{2i\pi \langle \cdot, \hat{r} \rangle}$ obtaining

$$h(t, z) = \frac{2}{\sqrt{3}} \sum_{\hat{r} \in \hat{R}} e^{2i\pi \langle z, \hat{r} \rangle} e^{-2\pi^2 |\hat{r}|^2 t}$$

Let us now integrate with respect to y in the triangle to obtain the distribution function for T_Ω , the first exit time of Brownian motion starting from x_0 ,

$$P(T_\Omega > t) = \int_{\Delta} p_t^0(x_0, y) dy.$$

We use the coordinates $y = y_1 + y_2 j$, $1 \geq y_1 \geq y_2 \geq 0$ on the triangle, with Jacobian $\frac{\sqrt{3}}{2}$ and $\hat{r} = ma + nb$ on $\hat{R}$, then

$$\langle y, \hat{r} \rangle = my_1 + ny_2, \quad \langle x_0, \hat{r} \rangle = 2m/3 + n/3, \quad |\hat{w}|^2 = 4(m^2 + mn + n^2)/3.$$

Furthermore, by periodicity

$$\int_{\Delta} h(t, x - \bar{y}) dy = \int_{\Delta} h(t, x - j - \bar{y}) dy = \int_{\bar{\Delta}+j} h(t, x - y) dy$$

and the triangle $\bar{\Delta} + j$ corresponds to the coordinates $1 \geq y_2 \geq y_1 \geq 0$. Let us compute

$$\int_{\Delta} h(t, x_0 - y) dy = \frac{\sqrt{3}}{2} \sum_{n,m} e^{2i\pi(2m/3+n/3)} \int_0^1 dy_1 \int_0^{y_1} e^{-2i\pi(my_1+ny_2)} dy_2 e^{-8\pi^2(m^2+mn+n^2)t/3}$$

It is easy to see that terms with $n \neq 0, m \neq 0, m+n \neq 0$ give 0 contribution. Also the term $m = n = 0$ will be cancelled by the other integral. The sequence $m = 0, n \neq 0$, gives

$$\begin{aligned} & \sum_{n \neq 0} e^{2i\pi n/3} \int_0^1 dy_1 \int_0^{y_1} e^{-2i\pi ny_2} dy_2 e^{-8\pi^2 n^2 t/3} \\ &= \sum_{n \neq 0} e^{2i\pi n/3} \int_0^1 \frac{1}{2i\pi n} (1 - e^{-2i\pi ny_1}) dy_1 e^{-8\pi^2 n^2 t/3} \\ &= \sum_{n \neq 0} e^{2i\pi n/3} \frac{1}{2i\pi n} e^{-8\pi^2 n^2 t/3} \end{aligned}$$

The sequence $m \neq 0, n = 0$, gives

$$\begin{aligned} & \sum_{m \neq 0} e^{4i\pi m/3} \int_0^1 dy_1 e^{-2i\pi my_1} \int_0^{y_1} dy_2 e^{-8\pi^2 m^2 t/3} \\ &= \sum_{m \neq 0} e^{4i\pi m/3} \int_0^1 dy_1 y_1 e^{-2i\pi y_1 m} e^{-8\pi^2 m^2 t/3} \\ &= - \sum_{m \neq 0} e^{4i\pi m/3} \frac{1}{2i\pi m} e^{-8\pi^2 m^2 t/3} \end{aligned}$$

and the sequence $m \neq 0, m+n=0$, gives

$$\begin{aligned} & \sum_{m \neq 0} e^{2i\pi m/3} \int_0^1 dy_1 \int_0^{y_1} e^{2i\pi(-my_1+my_2)} dy_2 e^{-8\pi^2 m^2 t/3} \\ &= \sum_{m \neq 0} e^{2i\pi m/3} \frac{1}{2i\pi m} \int_0^1 dy_1 (1 - e^{-2i\pi y_1 m}) e^{-8\pi^2 m^2 t/3} \\ &= \sum_{m \neq 0} e^{2i\pi m/3} \frac{1}{2i\pi m} e^{-8\pi^2 m^2 t/3} \end{aligned}$$

The sum of these three terms is

$$3 \sum_{n=1}^{\infty} \frac{\sin(2\pi n/3)}{\pi n} e^{-8\pi^2 n^2 t/3}.$$

The other integral is

$$\int_{\Delta} h(t, x_0 - \bar{y}) dy = \sum_{n,m} e^{2i\pi(2m/3+n/3)} \int_0^1 dy_2 \int_0^{y_2} e^{-2i\pi(my_1+ny_2)} dy_1 e^{-8\pi^2(m^2+mn+n^2)t/3}$$

Using

$$\int_0^1 dy_1 \int_0^1 e^{-2i\pi(my_1+ny_2)} dy_1 dy_2 = 0$$

if m or n is $\neq 0$ we see that, except for the term $m = n = 0$ the other terms are the opposite of what we computed, therefore after some rearrangement

$$P(T_{\Omega} > t) = 3\sqrt{3} \sum_{n=1}^{\infty} \frac{\chi_3(n)}{\pi n} e^{-8\pi^2 n^2 t/3}$$

where

$$\chi_3(n) = \frac{2}{\sqrt{3}} \sin(2\pi n/3)$$

is the multiplicative Dirichlet character modulo 3. The density is

$$P(T_{\Omega} \in dt) = 8\sqrt{3} \sum_{n=1}^{\infty} \pi n \chi_3(n) e^{-8\pi^2 n^2 t/3}.$$

II.4.1 A curious identity in law

Let S be the first hitting time of $3a$ by a three-dimensional Bessel process starting from a . The Laplace transform is

$$E[e^{-\lambda^2 S/2}] = \frac{3 \sinh(\lambda a)}{\sinh(3\lambda a)} = 3 \sum_{n=1}^{\infty} \chi_3(n) e^{-2n\lambda a}.$$

Inverting term by term gives the density

$$\sum_{n=0}^{\infty} 6an \frac{\chi_3(n) e^{-2n^2 a^2/t}}{\sqrt{2\pi t^3}}$$

We can also use the alternative expression

$$\frac{3 \sinh(\lambda a)}{\sinh(3\lambda a)} = \sum_{n=1}^{\infty} \frac{3\sqrt{3}\pi n \chi_3(n)}{\pi^2 n^2 + 9\lambda^2 a^2}$$

and invert term by term the Laplace transform to get

$$P(S \in dt) = \sum_{n=1}^{\infty} \frac{3\sqrt{3}\pi n \chi_3(n)}{18a^2} e^{-\pi^2 n^2 t / (18a^2)}$$

The agreement between these two expressions is an instance of the functional equation of theta series. Putting $8/3 = 1/(18a^2)$, or $a = \frac{1}{4\sqrt{3}}$ we see that the exit time T_Ω and the time S have the same distribution. Furthermore, the Mellin transform of this distribution is

$$E(T_\Omega^s) = \pi^{-2s} 8^{1-s} 3^{s+1/2} \Gamma(s+1) L_{\chi_3}(2s+1)$$

where L_{χ_3} is the Dirichlet L-function associated with the character χ_3 . Coming back to T_Ω , the triangle P_Ω has sides of size $2\sqrt{3}$ therefore

$$E[e^{-\frac{\lambda^2}{2} T_\Omega}] = \frac{3 \sinh(\lambda/2)}{\sinh(3\lambda/2)}.$$

Using (2.1) we are reduced to an elementary integral, and the result is

$$\ell_\Omega = \frac{\pi}{\sqrt{3}}.$$

II.5 The other cases

II.5.1 $\Omega = \{0, \pi/2\}$

We have

$$h_\Omega(\theta) = \max\{X(s) \cos \theta + Y(s) \sin \theta, Y(s) \cos \theta - X(s) \sin \theta ; 0 \leq s \leq t\}.$$

The trick is to use the fact that $(X(s) \cos \theta + Y(s) \sin \theta)_{s \geq 0}$ and $(Y(s) \cos \theta - X(s) \sin \theta)_{s \geq 0}$ are two standard one dimensional Brownian motions which are independent. As a consequence, using (2.3) we have

$$\Pr(h_\Omega(\theta) \leq h) = H^2(h/\sqrt{t})$$

where

$$H(z) = -1 + 2 \int_{-\infty}^z e^{-\frac{x^2}{2}} \frac{dx}{\sqrt{2\pi}}. \quad (2.8)$$

It leads to

$$\mathbb{E}(L(\Omega)) = 2\pi \int_0^\infty (1 - H^2(h/\sqrt{t})) dh = 2\pi\sqrt{t} \int_0^\infty (1 - H^2(z)) dz.$$

We write

$$1 - H^2(z) = 4 \int_z^\infty e^{-\frac{x^2}{2}} \frac{dx}{\sqrt{2\pi}} \left(1 - \int_z^\infty e^{-\frac{y^2}{2}} \frac{dy}{\sqrt{2\pi}} \right)$$

By the change of variables $x = zu$ and $y = zv$ and $w = z^2/2$ we get easily

$$\mathbb{E}(L(K_2(t))) = \sqrt{8\pi t} \left(2 - \int_1^\infty \int_1^\infty \frac{dudv}{(u^2 + v^2)^{3/2}} \right)$$

Introduce the function $(x, y) \mapsto g(x, y)$ defined for x and y in $\mathbb{R} \setminus \{0\}$ by

$$g(x, y) = \frac{\sqrt{x^2 + y^2}}{xy}. \quad (2.9)$$

A tedious calculation shows that if $a < b$ and $c < d$ with $(0, 0) \notin [a, b] \times [c, d]$ and if furthermore $a, b, c, d \neq 0$ we have

$$\int_a^b \int_c^d \frac{dudv}{(u^2 + v^2)^{3/2}} = g(a, c) + g(b, d) - g(a, d) - g(b, c). \quad (2.10)$$

From this we get that $\int_1^\infty \int_1^\infty \frac{dudv}{(u^2 + v^2)^{3/2}} = 2 - \sqrt{2}$ which finally gives that

$$\ell_\Omega = \sqrt{2}.$$

II.5.2 $\Omega = \{0, \pi/2, \pi, 3\pi/2\}$

The method is quite similar to the method used before, but the distribution function H of the maximum of a Brownian motion is replaced by the distribution function L of the maximum of its absolute value, which can be obtained from (2.6). Still with the previous notations we have

$$h_{C_\Omega}(\theta) = \max\{|X(s) \cos \theta + Y(s) \sin \theta|, |Y(s) \cos \theta - X(s) \sin \theta| ; 0 \leq s \leq t\}.$$

We use again the fact that $(X(s) \cos \theta + Y(s) \sin \theta)_{s \geq 0}$ and $(Y(s) \cos \theta - X(s) \sin \theta)_{s \geq 0}$ are two standard one dimensional independent Brownian motions. We get

$$\Pr(h_{C_\Omega}(\theta) \leq h) = L^2(h/\sqrt{t})$$

and

$$\mathbb{E}(L(C_\Omega(t))) = \int_0^{2\pi} \mathbb{E}(h_{C_4(t)}(\theta)) d\theta = 2\pi \int_0^\infty (1 - L^2(h/\sqrt{t})) dh = 2\pi\sqrt{t} \int_0^\infty (1 - L^2(z)) dz.$$

We now write

$$1 - L^2(z) = (1 - L(z))(1 + L(z)) = \frac{2}{\pi} \sum_{k=-\infty}^\infty \sum_{n=-\infty}^\infty \int_{(4k-1)z}^{(4k+1)z} e^{-\frac{x^2}{2}} dx \int_{(4n+1)z}^{(4n+3)z} e^{-\frac{y^2}{2}} dy.$$

We are led to the computation of the integral for fixed $(k, n) \in \mathbb{Z}^2$

$$I(k, n) = \sqrt{\frac{2}{\pi}} \int_0^\infty \left(\int_{(4k-1)z}^{(4k+1)z} e^{-\frac{x^2}{2}} dx \int_{(4n+1)z}^{(4n+3)z} e^{-\frac{y^2}{2}} dy \right) dz.$$

By the same change of variables as before $x = zu$ and $y = zv$ and $w = z^2/2$ we get

$$I(k, n) = \int_{4k-1}^{4k+1} \int_{4n+1}^{4n+3} \frac{dudv}{(u^2 + v^2)^{3/2}}. \quad (2.11)$$

In order to compute $I(k, n)$ we use the function g defined by (2.9) and the identity (2.10). From (2.11) it is apparent that $I(k, n) > 0$. To see also that

$$\ell_\Omega = \sum_{k=-\infty}^{\infty} \sum_{n=-\infty}^{\infty} I(k, n) < \infty \quad (2.12)$$

we use (2.11) again for observing that ℓ_Ω is the integral of the function $(u^2 + v^2)^{-3/2}$ over the union of squares

$$D = \cup_{k=-\infty}^{\infty} \cup_{n=-\infty}^{\infty} [4k-1, 4k+1] \times [4n+1, 4n+3]$$

which satisfies $D \subset A = \{(u, v) ; u^2 + v^2 \geq 1\}$. This implies, using polar coordinates $u = r \cos \theta$ and $v = r \sin \theta$:

$$\ell_\Omega = \int_D \frac{dudv}{(u^2 + v^2)^{3/2}} < \int_A \frac{dudv}{(u^2 + v^2)^{3/2}} = 2\pi \int_1^\infty r^{-2} dr = 2\pi.$$

In order to have some numerical evaluation of ℓ_Ω one can observe that we have the following two symmetries: for all $(k, n) \in \mathbb{Z}^2$

$$I(k, n) = I(-k, n) = I(k, -n-1).$$

This implies that ℓ_Ω can be written as $\ell_\Omega = S_0 + S_1$ with

$$S_0 = \sum_{n=-\infty}^{\infty} I(0, n) = 4 \sum_{n=0}^{\infty} \left(\frac{\sqrt{1 + (4n+1)^2}}{4n+1} - \frac{\sqrt{1 + (4n+3)^2}}{4n+3} \right) = 4 \sum_{m=0}^{\infty} (-1)^m u_m \quad (2.13)$$

with $u_m = \sqrt{1 + \frac{1}{(2m+1)^2}} - 1$, and $S_1 = 4 \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} I(k, n)$. Since $(u_m)_{m \geq 0}$ is decreasing we can estimate $\frac{S_0}{4} \approx 0.3725...$ We compute ℓ_Ω by Mathematica (we thank Daoud Bshouty for this) and we get

$$\ell_\Omega = 1,9178...$$

II.5.3 $\Omega = \{0, \pi/2, \pi\}$

Again the same method gives

$$\mathbb{E}(L(C_\Omega)) = 2\pi \int_0^\infty (1 - L(h/\sqrt{t})H(h/\sqrt{t}))dh = 2\pi\sqrt{t} \int_0^\infty (1 - L(z)H(z))dz.$$

We write for simplicity $L = 1 - 2L_1$ and $H = 1 - 2H_1$ so that

$$\mathbb{E}(L(C_\Omega)) = 2\pi\sqrt{t} \int_0^\infty (2L_1(z) + 2H_1(z) - 4L_1(z)H_1(z)) dz.$$

A previous calculation has shown that $2\pi\sqrt{t} \int_0^\infty 2L_1(z)dz = \frac{\pi}{2}\sqrt{8\pi t}$ and Letac (1978) -or a direct calculation- shows $2\pi\sqrt{t} \int_0^\infty 2H_1(z)dz = \sqrt{8\pi t}$. The delicate remaining point is the computation of $\int_0^\infty 4L_1(z)H_1(z)dz$. By the same methods as before, the same changes of variable $x = zu$ and $y = zv$ and $w = z^2/2$, the use of the function g of (2.9) and the formula (2.10) we get

$$\begin{aligned} 2\pi\sqrt{t} \int_0^\infty 4L_1(z)H_1(z)dz &= \sqrt{8\pi t} \sum_{n=-\infty}^{\infty} \int_1^\infty \int_{4n+1}^{4n+3} \frac{dudv}{(u^2 + v^2)^{3/2}} \\ &= \sqrt{8\pi t} \sum_{n=-\infty}^{\infty} (g(1, 4n+1) - g(1, 4n+3)). \end{aligned}$$

It is easily seen that with the notation (2.13)

$$\sum_{n=-\infty}^{\infty} (g(1, 4n+1) - g(1, 4n+3)) = \frac{S_0}{2} \approx 0.7450... \quad (2.14)$$

Thus

$$\ell_\Omega \approx \frac{\pi}{2} + 0.2550....$$

III Comments

1. The explicit calculations of the paper for the mean perimeter are feasible since we are able to have the exact distribution of the support functions $\theta \mapsto h_C(\theta)$ for the random convex sets considered here. The general case seems a difficult problem.
2. Calculations for the mean areas rely on the formula $A(C) = \int_0^{2\pi} (h_C(\theta)^2 - h'_C(\theta)^2) d\theta$, where $h'_C(\theta)$ denotes the derivative on the left, which can be shown to exist (see Letac (1983)) and ultimately of the distribution of $h'_C(\theta)$ when C is random. This problem was solved in the unpublished thesis of El Bachir (1983) for $C_1(t)$, where $\mathbb{E}(A(C_1(t))) = \pi t/2$ is proved. It is unsolved for the other convex sets considered here, except for $C_\infty(t)$ the smallest circle, centered at 0, surrounding $C_1(t)$, indeed

$$\mathbb{E}(A(C_\infty(t))) = \pi t \int_0^\infty \frac{ds}{I_0(\sqrt{2s})} = \pi t \times 3.06883.$$

a little more than six times the quantity $\mathbb{E}(A(C_1(t)))$. Thanks to Daoud Bshouty for the numerical evaluation of this integral.

IV References

BIANE, P.; PITMAN, J.; YOR, M. (2001) 'Probability laws related to the Jacobi theta and Riemann zeta functions, and Brownian excursions' *Bull. Amer. Math. Soc. (N.S.)* **38**, no. 4, 435–465.

DOUMERC, Y.: O'CONNELL, N (2001) 'Exit problems associated with finite reflection groups' *Probab. Th. Rel. Fields*) **132**, no. 4, 501–538.

EL BACHIR, M. (1983) *L'enveloppe convexe du mouvement brownien*. Thèse de 3ème cycle, Université Paul Sabatier, Toulouse.

FELLER, W. (1966) *An Introduction to Probability Theory and Its Applications*, Vol 2. Wiley, New York.

LETAC, G. (1978) 'Advanced problem 6230' *Am. Math. Monthly*. Solution by L. Takács **87**, 142 (1980).

LETAC, G. (1983) 'Mesures sur le cercle et convexes du plan' *Annales scientifiques de l'Université de Clermont-Ferrand II* **76**, 35-65.

LÉVY, P. (1948) *Processus stochastiques et mouvement brownien*, Gauthier Villars, Paris.